

### **Street Address:**

3356 Big Pine Trail, Ste C&D Champaign, IL 61822 USA **Phone:** +1-217-359-2344

E-Mail: CRM@aocs.org; Web: www.aocs.org

# Certified Reference Materials AOCS 0804-B2

Report of the certification process for

MON 1445 Cotton

**Certified Reference Materials** 

Second Batch

OECD Unique ID MON-Ø1445-2

Denise Williams
Technical Services Manager

Tiffanie West Technical Services Director



### **Legal Notice**

Neither AOCS nor any person acting on behalf of AOCS is responsible for the use which might be made of the following information.

### **AOCS Mission Statement**

AOCS advances the science and technology of oils, fats, proteins, surfactants and related materials, enriching the lives of people everywhere.

More information regarding AOCS is available at <a href="http://www.aocs.org">http://www.aocs.org</a>

# **Table of Contents**

Abstract	4
Acknowledgments	5
Glossary	6
Introduction	8
Material Processing	8
Trait Verification to Certify Presence of MON 1445	8
Certified Value and Measurement Uncertainty	9
Homogeneity	10
Stability	10
References	13

### **Abstract**

This report describes the preparation and certification of the cotton CRM AOCS 0804-B2 produced by AOCS Technical Services in 2020. The CRMs have been prepared according to ISO 17034:2016 and are intended to serve as control material for third party testing of cottonseeds for transformation events. The cotton MON 1445 powder was provided by Bayer CropScience, St. Louis, MO (hereinafter "Bayer CropScience"). It was prepared by grinding the bulk seed at Bayer CropScience. The certified value of AOCS 0804-B2 was based on the purity of the bulk seed material and with 95% confidence, the true value is ≥ 975 g/kg. The powder was aliquoted and bottled in 27-mL glass headspace vials and sealed under a nitrogen gas environment at Illinois Crop Improvement Association. The presence of MON 1445 in AOCS 0804-B2 was verified using event-specific, qualitative PCR analysis by Eurofins-GeneScan, New Orleans, LA (an ISO 17025 accredited laboratory). CRM samples should be stored dry in a sealed container at ambient or cooler conditions in the dark.

## **Acknowledgements**

The authors would like to express sincere appreciation and gratitude to several individuals and their companies for support and guidance throughout this project. Thanks go to Jack Milligan, Bayer CropScience, for offering AOCS the opportunity to manufacture and distribute these products; to Sandra Harrison, Charlie Drennan and the crew at Illinois Crop Improvement Association for packaging the samples; and to Frank Spiegelhalter, Greg Ditta, E. Pearce Smith, and Daniel Thompson, Eurofins-GeneScan for event-specific, qualitative PCR analysis including the provision of information on running the analyses and interpreting the results.

### Glossary

AOCS American Oil Chemists' Society

Conventional Crop Crop variety with no history of modern biotechnology and is

produced through plant-breeding techniques that rely on selecting and mating parent plants possessing promising traits and repeatedly selecting for superior performance

among their offspring

DNA Deoxyribonucleic Acid

Detection Limit Lowest level at which target DNA can be detected in a sample

and be reliably tested by PCR methods. It is typically expressed as a percentage: the ratio of the number of modern

biotechnology derived genomes to the number of crop

genomes times 100 percent

EC European Commission

Genome The full set of genes and associated DNA characteristic of an

organism

GMO Organism that has had genetic sequences modified using

molecular-level techniques

ISO International Organisation for Standardisation

PCR Polymerase Chain Reaction: technique used to determine

whether a sample of plant tissue contains a particular DNA sequence. PCR relies on primer sets that zero in on a

particular target DNA sequence and a special DNA-copying

enzyme (DNA polymerase) that makes enough copies of the

target sequence for identification and measurement

Report of Certification for 0804-B2 Page 6 of 13 ©AOCS, 2024 Qualitative PCR PCR methods that determine the presence or absence of a

specific target DNA sequence at a particular level of detection

Quantitation Limit Lowest level at which the amount of target DNA sequence in

a sample can be reliably quantified. It is typically expressed

as the ration of the number of transgenic genomes to the

number of crop genomes times 100 percent.

Quantitative PCR PCR methods that estimate the relative amount of target DNA

sequence in a mixture of DNA molecules

Trait: MON 1445 Roundup Ready® Cotton

Introduction

Plant genetic modification is an extension of traditional plant breeding. It allows plant

breeders to develop crops with specific traits including insect, disease, and herbicide

resistance; processing advantages; and nutritional enhancement. An important

component for identifying these new traits is a Certified Reference Material created from

leaf, seed, or grain containing the new trait as well as a CRM created from the

conventionally bred matrix. The European Commission has mandated that from 18 April

2004, a method for detecting a new event derived from modern biotechnology and

Certified Reference Material must be available before the EC will consider authorizing

acceptance of a new crop derived from modern biotechnology. Several nations outside

Europe also require grain and ingredients to be labeled above a threshold before

accepting a shipment.

To meet the above analytical requirements for GM determination, AOCS 0804-B2 was

manufactured from cottonseeds according to ISO 17034:2016 and in accordance with EC

No 1829/2003, EC No 641/2004 and EC No 619/2011. This CRM is available from AOCS.

**Material Processing** 

The MON 1445 cotton seed used to prepare AOCS 0804-B2 were homozygous resulting

from several cycles of self-pollination. Bayer CropScience milled ~4 kg of MON 1445

cotton seed. All the seed powder was passed through a 710 µm mesh sieve. The seed

powder was delivered to AOCS who contracted Illinois Crop Improvement Association for

packaging the samples. The powder was aliquoted and bottled in 27-mL glass headspace

vials and sealed under a nitrogen gas environment.

**Trait Verification to Certify Presence of MON 1445** 

The presence of the MON 1445 trait was assessed on 10 random vials of AOCS 0804-B2. AOCS used the Random Number Generator function of Microsoft Excel to select samples for verification of trait presence. Sample numbers that were randomly selected were sent to Eurofins-GeneScan, New Orleans, LA (an ISO 17025 accredited laboratory) for event-specific, qualitative PCR analysis to verify the presence of MON 1445 in the samples (Table 1).

Table 1. Trait verification testing on AOCS 0804-B2 MON 1445 maize performed by Eurofins-GeneScan, New Orleans, LA (an ISO 17025 accredited laboratory).

AOCS 0804-B2 Sample	Trait MON 1445 Presence	
Sample # 20	Positive	
Sample # 51	Positive	
Sample # 107	Positive	
Sample # 144	Positive	
Sample # 177	Positive	
Sample # 252	Positive	
Sample # 267	Positive	
Sample # 344	Positive	
Sample # 379	Positive	
Sample # 384	Positive	

# **Certified Value and Measurement Uncertainty**

The genetic purity of the seed lot used to produce AOCS 0804-B2 was assessed by Bayer CropScience. A total of 704 cottonseed seeds were subjected to individual seed testing for the presence of MON 1445 by qualitative event-specific PCR. 702 of the 704 seeds tested positive for the presence of MON 1445.

Purity estimation was calculated using SeedCalc8 (Remund *et al.*, 2008) and corresponds to the lower bound of true % purity. The % purity in the sample was 99.7%, when 704

seeds were tested. Using a 95% confidence level, the true % purity of the MON 1445 seed lot was at least 97.5%. Consequently, with 95% confidence, the true value is ≥ 975 g/kg.

The Measurement Uncertainty was based on the lower bound of the true % purity and is the expanded uncertainty with a coverage factor of 2 and confidence level of 95%. It is obtained by combining the uncertainties from the purity assessment  $(u_{char,rel})$ , the homogeneity assessment  $(u_{bb,rel})$ , the transport stability assessment  $(u_{sts,rel})$  and the long-term stability assessment  $(u_{lts,rel})$ :

$$u_{CRM,rel} = \sqrt{u_{char,rel}^2 + u_{bb,rel}^2 + u_{sts,rel}^2 + u_{lts,rel}^2}$$

$$U_{CRM} = 2 \times u_{CRM,rel} \times 1000 \ g/kg$$

When using an asymmetric uncertainty, the reported measurement uncertainty is truncated on the right side such that the value does not exceed 1000 g/kg. Consequently, the expanded measurement uncertainty for AOCS 0804-B2 is +3 g/kg, -23 g/kg.

# Homogeneity

The homogeneity of AOCS 0804-B2 is related to the purity of the seeds. 702 out of 704 cottonseed seeds tested positive for the MON 1445 cottonseed event by event-specific PCR. Based on the sample purity of 99.7%, as determined using SeedCalc8, the batch was considered to be homogenous.

In addition, the homogeneity of the MON 1445 trait was confirmed when 10 random vials of AOCS 0804-B2 were selected and were sent to Eurofins-GeneScan, New Orleans, LA (an ISO 17025 accredited laboratory) for event-specific, qualitative PCR analysis to verify the presence of MON 1445 in the samples (See Trait Verification section and Table 1).

# **Stability**

Time, temperature and light are regarded as the most relevant influences on the stability

of CRM (Linsinger, et al., 2001). The influence of light is mitigated by shipping and storing

the vials in boxes, thus minimizing the possibility of degradation due to light. The influence

of temperature is mitigated by storing the vials in a temperature-controlled room, and

shipping vials at ambient temperature.

The effect of temperature and time are investigated.

A transport (short-term) stability study is conducted to assess the stability of maize CRM

during transport. The temperature and time conditions in the study cover the typical

conditions and the not so rare situations. The outcome of the study is considered

transferable to other CRMs of similar property. Samples were subject to 3 different

temperatures (4 °C (fridge), 25 °C (ambient), 60 °C (oven)) for 4 different durations (0, 1,

2, and 4 weeks). The study concluded that samples are stable at 4 °C (fridge) and 25 °C

(ambient) for 4 weeks. The estimated uncertainty contribution from transport (short-term)

stability is 1.0%.

A long-term stability study is conducted to assess the stability of maize CRM during

storage. Samples are stored at 25 °C (ambient) and the stability of the sample is monitored

as long as the samples is available. The storage temperate studied is 25 °C and the length

of time to be studied is 10 years. The outcome of the study is considered transferable to

other CRMs of similar property. In the initial 1-year stability study, samples were subject

the storage condition for 4 different durations (0, 1, 3, 6 and 12 months). The study

concluded that samples are stable at 25 °C (ambient) for 12 months. The estimated

uncertainty contribution from long-term stability is 0.42%.

Stability of these CRMs has been listed as 1 year from the certification date. The

materials were processed and are stored at ambient or lower temperature, under

nitrogen, in glass headspace vials. These materials are expected to be stable for longer

than the estimated expiration date. The stability of the powder material will be reevaluated

Report of Certification for 0804-B2

at time of expiration.		e still representativ	ve of the certified	value, the
certificates will be exte	ended.			

### References

Eurofins-GeneScan; 2219 Lakeshore Drive, Suite 400, New Orleans, LA 70122; Telephone: +1 504 297 4330 Toll Free: +1 866 535 2730 Fax: +1 504 297 4335 <a href="https://www.eurofinsus.com/food-testing/testing-services/gmo/">https://www.eurofinsus.com/food-testing/testing-services/gmo/</a>

Illinois Crop Improvement Association, 3105 Research Road, Champaign, IL 61826; Telephone: +1 217 359 4053 Fax: +1 217 359 4075; <a href="http://www.ilcrop.com/index.htm">http://www.ilcrop.com/index.htm</a>

ISO 17034:2016 (E) General requirements for the competence of reference material producers

ISO 17025:2005 and ISO 17025:2017, General Requirements for the Competence of Testing and Calibration Laboratories

International Seed Testing Association, International Rules of Seed Testing: Seed Science and Technology Rules, 2012

Regulation (EC) No 1829/2003 of the European Parliament and of the Council of 22 September 2003 on genetically modified food and feed; <a href="https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX%3A32003R1829&amp;from=en">https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX%3A32003R1829&amp;from=en</a>

Remund K., Simpson R., Laffont J-L., Wright D., and Gregoire S. Seedcalc8. 2008. <a href="https://www.seedtest.org/en/statistical-tools-for-seed-testing-content---1-3449--1102.html">https://www.seedtest.org/en/statistical-tools-for-seed-testing-content---1-3449--1102.html</a>